

## APPLE SPRAY PROGRAM

USE	AUGUST	SEPTEMBER	OCTOBER	NOVEMBER	DECEMBER	JANUARY	FEBRUARY	MARCH	APRIL	MAY	
<b>Etherics 1000</b> General	1 x per month										
<b>ThermoMax</b> Frost / Warming		3 weekly									
<b>PhotoMax</b> Photosynthesis		with ThermoMax									
<b>FungalMin</b> Fungal control		every two weeks									
<b>SilicaMax</b> Hardening				once		once		once			
<b>Etherics 7</b> Slow ripening							2x a week apart 1 month before harvest				
<b>RipeMax</b> Increase ripening							2 x 2 weeks apart 1 month before harvest				
<b>Zeroln</b> Reduce water						2 x 1 week apart 3 weeks before harvest					

### ThermoMax

ThermoMax is to be applied every 3 weeks during spring time, for control of frost.

It has a warming effect and so can be continued to be used through till December, if the cold weather persists.

### Silica Max

SilicaMax brings a inner strength to plants. It can be sprayed 2 weeks after fruit set, once in January, if there is not a drought, and 3 weeks before harvest.

### Etherics 1000

Can be used monthly with liquid fertilisers, otherwise every 2nd month

### FungalMin - Trial

This is to be sprayed every 2 weeks throughout the whole season

### PhotoMax

PhotoMax is used to enhance photosynthesis within the plant. This is very useful in early spring and if dull weather persists throughout the season. Higher dry matter has been experienced where it is used monthly throughout the season.

### Zeroln

Zeroln is used to reduce the water uptake of a plant after rain. Best to use on stone fruit like plums, cherries, apricots and peaches etc.

If it looks like a heavy rain period is likely over harvest. Spray once a two weeks before harvest and a second spray just before the rain.

If the rain is heavy add SilicaMax to the second spray. Once Zeroln has been applied fruit sizing will slow down.

### Etherics 7- Trial

Etherics 7 can be used to slow the ripening process. Apply once a month prior to harvest and again two weeks later.

Fruit size should continue to grow. If ripening is then needed to be sped up use RipeMax

### RipeMax - Trial

RipeMax can be used to sped up ripening. Apply once a month before harvest and then a second spray two weeks before harvest.



More Product  
information can  
be found at  
[www.bdmax.co.nz](http://www.bdmax.co.nz).  
Or phone  
07 573 5859  
or 0800735859